

On Ritt's decomposition Theorem in the case of finite fields

Jaime Gutierrez*

Dpto. de Matemáticas, Estadística y Computación, Universidad de Cantabria
E-39071 Santander, Spain
jaime.gutierrez@unican.es

David Sevilla

Department of Computer Science, Concordia University
Montreal H3G 1M8, QC, Canada
dsevilla@cs.concordia.ca

Abstract

A classical theorem by Ritt states that all the complete decomposition chains of a univariate polynomial satisfying a certain tameness condition have the same length. In this paper we present our conclusions about the generalization of these theorem in the case of finite coefficient fields when the tameness condition is dropped. (*Updated April 2008: see note at the beginning of the introduction.*)

1 Introduction

(Updated April 2008: There exists a previously published article, of which the authors were not aware, where Ritt's second theorem is shown to be true in all characteristics, provided we have the natural and necessary condition that no derivative vanishes identically. This goes much beyond the condition

*Both authors are partially supported by Research Project MTM2004-07086 of the Spanish Ministry of Science and Technology

that p does not divide the degrees (which is not a necessary condition). See additional references [18] and [19]. We wish to thank Pr. Zannier for this information.)

Our starting point is the decomposition of polynomials and rational functions in one variable. First we define the basic concepts of this topic.

Definition 1. Let $\mathbb{K}$ be any field, x a transcendental over $\mathbb{K}$ and $\mathbb{K}(x)$ the field of rational functions in the variable x with coefficients in $\mathbb{K}$. In the set $T = \mathbb{K}(x) \setminus \mathbb{K}$ we define the binary operation of *composition* as

$$g(x) \circ h(x) = g(h(x)) = g(h).$$

We have that $(T, \circ)$ is a semigroup, the element x being its neutral element.

If $f = g \circ h$, we call this a *decomposition* of f and say that g is a *component on the left* of f and h is a *component on the right* of f . We call a decomposition *trivial* if any of the components is a unit with respect to decomposition.

Given two decompositions $f = g_1 \circ h_1 = g_2 \circ h_2$ of a rational function, we call them *equivalent* if there exists a unit u such that

$$h_1 = u \circ h_2 \quad (\text{thus, } g_1 = g_2 \circ u^{-1}),$$

where the inverse is taken with respect to composition.

Given $f \in T$, we say that it is *indecomposable* if it is not a unit and all its decompositions are trivial.

We define a *complete* decomposition of $f \in \mathbb{K}(x)$ to be $f = g_1 \circ \cdots \circ g_r$ where every g_i is indecomposable. The notion of equivalent complete decompositions is straightforward from the previous concepts.

Definition 2. Given a non-constant rational function $f(x) \in \mathbb{K}(x)$ where $f(x) = f_N(x)/f_D(x)$ with $f_N, f_D \in \mathbb{K}[x]$ and $(f_N, f_D) = 1$, we define the *degree* of f as

$$\deg f = \max\{\deg f_N, \deg f_D\}.$$

We also define $\deg a = 0$ when $a \in \mathbb{K}$.

From now on, we will use the previous notation when we refer to the numerator and denominator of a rational function. Unless explicitly stated, we will take the numerator to be monic, even though multiplication by constants will not be relevant.

Now we can properly state the problem of decomposition of univariate rational functions, although this will not be our main object of study.

Problem 3. Given a univariate rational function, decide if it is decomposable, and in the affirmative case compute a non-trivial decomposition of the function.

It is clear that the solution of this problem provides the computability of a complete decomposition of a function if it exists.

Next, we introduce some basic results about univariate decomposition, see [1] for more details.

Lemma 4.

- (i) For every $f \in T$, $\deg f = [\mathbb{K}(x) : \mathbb{K}(f)]$.
- (ii) $\deg(g \circ h) = \deg g \cdot \deg h$.
- (iii) $f(x)$ is a unit with respect to composition if and only if $\deg f = 1$, that is, $f(x) = \frac{ax+b}{cx+d}$ with $a, b, c, d \in \mathbb{K}$ and $ad - bc \neq 0$.
- (iv) Every non-constant element of $\mathbb{K}(x)$ is cancelable on the right with respect to composition. In other words, if $f(x), h(x) \in T$ are such that $f(x) = g(h(x))$ then $g(x)$ is uniquely determined by $f(x)$ and $h(x)$.

We can relate decomposition and Field Theory by means of the following classical result:

Theorem 5 (Lüroth's Theorem). *Let $\mathbb{F}$ be a field such that $\mathbb{K} \subset \mathbb{F} \subset \mathbb{K}(x)$. Then there exists $f \in \mathbb{K}(x)$ such that $\mathbb{F} = \mathbb{K}(f)$. Also, if $\mathbb{F}$ contains a polynomial, f can be chosen to be a polynomial.*

Proof. See for example [9] for a proof in the case $\mathbb{K} = \mathbb{C}$, [15] for one in the general case and [16] for an elementary one. Constructive proofs can be found in [10], [13] and [1]. □

Now we state one of the classical Ritt's theorems (see [11]) about the relations among the complete decompositions of a polynomial that satisfies a certain condition. First we have to define that condition.

Definition 6. A polynomial $f \in \mathbb{K}[x]$ is *tame* when $\text{char } \mathbb{K}$ does not divide $\deg f$.

Ritt's theorem essentially proves that all the decompositions have the same length and are related in a rather direct way.

Definition 7. A *bidecomposition* is a 4-tuple of polynomials f_1, g_1, f_2, g_2 such that $f_1 \circ g_1 = f_2 \circ g_2$, $(\deg f_1, \deg g_1) = 1$ and $\deg f_1 = \deg g_2$.

Theorem 8 (Ritt's Theorem). *Let $f \in \mathbb{K}[x]$ be tame and let $f = g_1 \circ \cdots \circ g_r = h_1 \circ \cdots \circ h_s$ be two complete decompositions of f . Then $r = s$, and the sequences $(\deg g_1, \dots, \deg g_r)$, $(\deg h_1, \dots, \deg h_s)$ are permutations of each other. Moreover, there exists a finite chain of complete decompositions*

$$f = f_1^{(j)} \circ \cdots \circ f_r^{(j)}, \quad j \in \{1, \dots, k\},$$

such that

$$f_i^{(1)} = g_i, \quad f_i^{(k)} = h_i, \quad i = 1, \dots, r,$$

and for each $j < k$, there exists i_j such that the j -th and $(j+1)$ -th decomposition differ only in one of these aspects:

- (i) $f_{i_j}^{(j)} \circ f_{i_j+1}^{(j)}$ and $f_{i_j}^{(j+1)} \circ f_{i_j+1}^{(j+1)}$ are equivalent.
- (ii) $f_{i_j}^{(j)} \circ f_{i_j+1}^{(j)} = f_{i_j}^{(j+1)} \circ f_{i_j+1}^{(j+1)}$ is a bidecomposition.

Proof. See [11] for $\mathbb{K} = \mathbb{C}$, [5] for characteristic zero fields and [6] for the general case. $\square$

In this paper we will study the generalization of this result to polynomials with coefficients in finite fields. To that end, we will also analyze the structure of intermediate fields between $\mathbb{K}(f)$ and $\mathbb{K}(x)$. It is already known that Ritt's theorem is false when the tameness condition is dropped, see [4] for a counterexample.

Let $f = g(h)$. Then $f \in \mathbb{K}(h)$, thus $\mathbb{K}(f) \subset \mathbb{K}(h)$. Also, $\mathbb{K}(f) = \mathbb{K}(h)$ if and only if $f = u \circ h$ for some unit u . This allows the following bijection among decompositions of a function f and fields between $\mathbb{K}(f)$ and $\mathbb{K}(x)$:

Theorem 9. *Let $f \in \mathbb{K}(x)$. In the set of decompositions of f we have the equivalence relation given by the definition of equivalence of decompositions. If we denote as $[(g, h)]$ the class of the decomposition $f = g(h)$, then we have the bijection:*

$$\begin{aligned} \{[(g, h)] : f = g(h)\} &\longleftrightarrow \{\mathbb{F} : \mathbb{K}(f) \subset \mathbb{F} \subset \mathbb{K}(x)\} \\ [(g, h)] &\longleftrightarrow \mathbb{F} = \mathbb{K}(h). \end{aligned}$$

Thanks to the Primitive Element Theorem (see for example [7]), we know that for each non-constant $f \in \mathbb{K}(x)$ there exist finitely many fields between $\mathbb{K}(f)$ and $\mathbb{K}(x)$. Due to the second part of Lüroth's Theorem, every rational decomposition of a polynomial is equivalent to a decomposition whose components are polynomials. Therefore it suffices to care about polynomial decomposition in this case.

In Section 2 we introduce several elementary results about univariate function fields that arise from Galois theory. In Section 3 we present a function that is fixed by all the automorphisms of a univariate function field over a finite field and several results related to it. In particular, we provide an essentially new counterexample of Ritt's theorem for finite coefficient fields.

2 The fixing group and the fixed field

In this section we introduce several simple notions from the classical Galois theory. Let $\Gamma(\mathbb{K}) = \text{Aut}_{\mathbb{K}}\mathbb{K}(x)$ (we will write simply Γ if there can be no confusion about the field). The elements of $\Gamma(\mathbb{K})$ can be identified with the images of x under the automorphisms, that is, with Möbius transformations (non-constant rational functions of the form $(ax+b)/(cx+d) \in \mathbb{K}(x)$), which are also the units of $\mathbb{K}(x)$ under composition.

Definition 10.

- Let $f \in \mathbb{K}(x)$. We define $G(f) = \{u \in \Gamma(\mathbb{K}) : f \circ u = f\}$.
- Let $H < \Gamma(\mathbb{K})$. We define $\text{Fix}(H) = \{f \in \mathbb{K}(x) : f \circ u = f \ \forall u \in H\}$.

This definitions correspond to the classical Galois correspondences (not bijective in general) between the intermediate fields of an extension and the subgroups of its automorphism group, as the following diagram shows:

$$\begin{array}{ccc}
 \mathbb{K}(x) & \longleftrightarrow & \{id\} \\
 | & & | \\
 \mathbb{K}(f) & \longrightarrow & G(f) \\
 | & & | \\
 \text{Fix}(H) & \longleftarrow & H \\
 | & & | \\
 \mathbb{K} & \longleftrightarrow & \Gamma
 \end{array}$$

Remark 11. As $\mathbb{K}(f) = \mathbb{K}(f')$ if and only if $f = u \circ f'$ for some unit u , we have that the application $\mathbb{K}(f) \mapsto G(f)$ is well-defined.

We are interested in the computability of these elements, the following results solves one of the two parts of this question.

Theorem 12. *Let $H = \{h_1, \dots, h_m\} \subset \mathbb{K}(x)$ be a finite subgroup of Γ . Let $P(T) = \prod_1^m (T - h_i) \in \mathbb{K}(x)[T]$. Then any non-constant coefficient of $P(T)$ generates $\text{Fix}(H)$.*

Proof. Sketch of proof. It can be shown that $P(T)$ is the minimal polynomial of x over $\text{Fix}(H) \subset \mathbb{K}(x)$. Then, a known proof of Lüroth's theorem (see [10]) gives the desired result. $\square$ $\square$

The previous theorem obviously provides an algorithm to compute the fixed field for a given finite subgroup of Γ : compute the symmetric elementary functions in $h_1, \dots, h_m$ until a non-constant one is found.

About the computation of the fixing group, an elementary but inefficient algorithm is given by the resolution of the equations given by

$$f(x) - f\left(\frac{ax+b}{cx+d}\right) = 0$$

in terms of a, b, c, d . Another algorithm (see [14]) combines this idea with certain normalization of the rational function, which simplifies the equations substantially.

Next, we state several interesting properties of the fixed field and the fixing group, see [14] for details.

Theorem 13. *Let $H < \Gamma$.*

- *H is infinite $\Rightarrow \text{Fix}(H) = \mathbb{K}$.*
- *H is finite $\Rightarrow \mathbb{K} \subsetneq \text{Fix}(H)$, $\text{Fix}(H) \subset \mathbb{K}(x)$ is a normal extension, and in particular $\text{Fix}(H) = \mathbb{K}(f)$ with $\deg f = |H|$.*

Theorem 14.

- (i) *Given a non-constant $f \in \mathbb{K}(x)$, $|G(f)|$ divides $\deg f$. Moreover, for any field $\mathbb{K}$ there is always a function $f \in \mathbb{K}(x)$ such that $1 < |G(f)| < \deg f$.*

- (ii) $|G(f)| = \deg f \Rightarrow \mathbb{K}(f) \subseteq \mathbb{K}(x)$ is normal. Moreover, if the extension $\mathbb{K}(f) \subseteq \mathbb{K}(x)$ is separable, then

$$\mathbb{K}(f) \subseteq \mathbb{K}(x) \text{ is normal} \Rightarrow |G(f)| = \deg f.$$

- (iii) Given a finite subgroup H of Γ , there is a bijection between the subgroups of H and the fields between $\text{Fix}(H)$ and $\mathbb{K}(x)$. Also, if $\text{Fix}(H) = \mathbb{K}(f)$, there is a bijection between the right components of f (up to equivalence by units) and the subgroups of H .

Proof. For the first item, we take $f = x^2(x-1)^2$ gives $G(f) = \{x, 1-x\}$. The other ones are straightforward. $\square$ $\square$

3 Finite fields

In this section, $\mathbb{K} = \mathbb{F}_q$ where $q = p^m$ and $p = \text{char } \mathbb{F}_q$, see [8] for several useful results. As before, we will denote $\Gamma = \Gamma(\mathbb{F}_q)$.

Definition 15. For any $\mathbb{K}$, $\Gamma_0 = \Gamma \cap \mathbb{K}[x] = \{ax + b : a \in \mathbb{K}^*, b \in \mathbb{K}\}$.

Theorem 16. $\mathbb{K}(x)$ is Galois over $\mathbb{K}$ (that is, the only functions fixed by $\Gamma(\mathbb{K})$ are the constants) if and only if $\mathbb{K}$ is infinite.

Proof. The "if" part is the first part of Theorem 13. The "only if" part is a consequence of Theorem 12, as $\Gamma(\mathbb{K})$ is finite whenever $\mathbb{K}$ is finite. $\square$

The interest of Γ and Γ_0 in the case of finite fields lies in the fact that both groups provide non-trivial fixed fields.

Theorem 17. The fixed field for Γ_0 is generated by $(x^q - x)^{q-1}$.

Proof. According to Theorem 12 any non-constant coefficient of $Q(T) = \prod_{u \in \Gamma_0} (T - u)$ generates the field. But the constant term of Q is precisely $\prod_{u \in \Gamma_0} u = (x^q - x)^{q-1}$. $\square$ $\square$

From now on, we will denote $P_q = (x^q - x)^{q-1}$.

As $\Gamma_0 \subset \Gamma$, if f generates the fixed field for Γ then $f = h(P_q)$ for some $h \in \mathbb{K}(x)$. Moreover, h has degree $[\Gamma : \Gamma_0] = q + 1$.

Theorem 18. *Let*

$$h_q = (x^{q+1} + x + 1)/x^q.$$

Then the rational function $f_q = h_q(P_q)$ generates $\text{Fix}(\Gamma)$.

Proof. It is easy to prove that $\Gamma_0 \cup \{1/x\}$ generates Γ . As f_q is a function of P_q and its degree is equal to the order of the group, it suffices to show that $f_q(1/x) = f_q(x)$. A simple computation shows that this is indeed the case: let $y = x^{q-1}$. Then $P_q(x) = y(y-1)^{q-1}$ and $P_q(1/x) = (y-1)^{q-1}/y^q$. Thus,

$$\begin{aligned} f_q(1/x) - f_q(x) &= \\ &= \frac{\frac{(y-1)^{q^2-1}}{y^{q^2+q}} + \frac{(y-1)^{q-1}}{y^q} + 1}{\frac{(y-1)^{q^2-q}}{y^{q^2}}} - \frac{y^{q+1}(y-1)^{q^2-1} + y(y-1)^{q-1} + 1}{y^q(y-1)^{q^2-q}} = \\ &= \frac{(y-1)^{q^2-1} + y^{q^2}(y-1)^{q-1} + y^{q^2+q} - y^{q+1}(y-1)^{q^2-1} - y(y-1)^{q-1} - 1}{y^q(y-1)^{q^2-q}} = \\ &= \frac{(y-1)^{q^2-1}(1 - y^{q+1}) + (y-1)^{q-1}(y^{q^2} - y) + y^{q^2+q} - 1}{y^q(y-1)^{q^2-q}} = \\ &= \frac{(y-1)^{q^2-1}(1 - y^{q+1}) + (y-1)^{q-1}((y-1)^{q^2} - (y-1)) + y^{q^2+q} - 1}{y^q(y-1)^{q^2-q}} = \\ &= \frac{(y-1)^{q^2-1}(1 - y^{q+1} + (y-1)^q) - (y-1)^q + y^{q^2+q} - 1}{y^q(y-1)^{q^2-q}} = \\ &= \frac{(y-1)^{q^2-1}(1 - y^{q+1} + y^q - 1) - (y-1)^q + (y^{q+1} - 1)^q}{y^q(y-1)^{q^2-q}} = \\ &= \frac{-(y-1)^{q^2}y^q - (y-1)^q + (y-1)^q(1 + y + \dots + y^q)^q}{y^q(y-1)^{q^2-q}} = \\ &= \frac{-(y-1)^{q^2}y^q + (y-1)^q(y + \dots + y^q)^q}{y^q(y-1)^{q^2-q}} = \\ &= \frac{-(y-1)^{q^2} + (y-1)^q(1 + \dots + y^{q-1})^q}{(y-1)^{q^2-q}} = \\ &= \frac{-(y-1)^{q^2} + (y^q - 1)^q}{(y-1)^{q^2-q}} = 0. \end{aligned}$$

□

□

Let $f \in \mathbb{F}_q(x)$. Let $\mathcal{C} = \{\mathbb{K} : \mathbb{F}_q \subseteq \mathbb{K} \subseteq \mathbb{F}_q(x)\}$ and

$$\begin{array}{ccc} \phi : & \mathcal{C} & \longrightarrow \\ & \mathbb{F}_q(f) & \longrightarrow \text{Fix}(G(f)) = \mathbb{F}_q(f') \end{array}$$

which is a well-defined application. Then it is easy to check that f' is a (not necessarily proper) right-component of f . Also, as $G(f) \subset \Gamma$, f' is a right-component of f_q . Thus, $\mathbb{F}_q(f) \subseteq \mathbb{F}_q(f')$ and $\mathbb{F}_q(f_q) \subseteq \mathbb{F}_q(f')$, therefore $\mathbb{F}_q(f, f_q) \subseteq \mathbb{F}_q(f')$.

Theorem 19. $\mathbb{F}_q(f, f_q) = \mathbb{F}_q(f')$.

Proof. Let $\mathbb{F}_q(f, f_q) = \mathbb{F}_q(m)$. Then there is a rational function $r(x, y)$ such that $r(f, f_q) = m$. For every $u \in G(f)$, $m \circ u = r(f \circ u, f_q \circ u) = r(f, f_q) = m$. Therefore, $m \in \text{Fix}(G(f)) = \mathbb{F}_q(f') \Rightarrow \mathbb{F}_q(m) \subseteq \mathbb{F}_q(f')$. The other part is straightforward. $\square$ $\square$

The polynomial P_q has at least two different decompositions:

$$P_q = x^{q-1} \circ (x^q - x) = (x(x-1)^{q-1}) \circ x^{q-1}.$$

This gives at least two decompositions for h_q , both involving the component $\frac{x^{q+1} + x + 1}{x^q}$.

Theorem 20.

- (i) $\frac{x^{q+1} + x + 1}{x^q}$ is indecomposable.
- (ii) $x^q - x$ is decomposable iff q is composite, that is, $q = p^m$ with $m \geq 2$.
- (iii) $x(x-1)^{q-1}$ is indecomposable.

Proof. (i) We will prove that for certain units $u, v \in \mathbb{F}_q(x)$, the function

$$u \circ \frac{x^{q+1} + x + 1}{x^q} \circ v$$

is indecomposable. In particular, let $u = x + 1, v = 1/(x - 1)$. Then

$$u \circ \frac{x^{q+1} + x + 1}{x^q} \circ v = \frac{x^{q+1}}{x - 1}.$$

As the degree is multiplicative with respect to composition, and so is the difference in the degrees of numerator and denominator (see [14, Theorem 1.14 and Corollary 1.15]), there is no possible decomposition for this function and the original function is also indecomposable.

- (ii) As $G(x^q - x) = \{x - a : a \in \mathbb{F}_q\}$ and $|G(x^q - x)| = q = \deg x^q - x$, by Theorem 14 there is a bijection between the decompositions of $x^q - x$ and the subgroups of its fixing group. But $G(x^q - x)$ has proper subgroups if and only if its order is composite.
- (iii) Let $q = p^m$. Let $x(x - 1)^{q-1} = g(h)$ with $g = x^{p^r} + g_0, \deg g_0 \leq p^r - 1$ and $h = x^{p^s} + h_0, \deg h_0 \leq p^s - 1$. Then

$$g \circ h = h^{p^r} + g_0 \circ h = (x^{p^s} + h_0)^{p^r} + g_0 \circ h = x^q + h_0^{p^r} + g_0 \circ h$$

with $\deg h_0^{p^r} \leq q - p^r$ and $\deg g_0 \circ h \leq q - p^s$. But

$$x(x - 1)^{q-1} = x^q + x^{q-1} + \dots + x^2 + x,$$

thus either $r = 0$ or $s = 0$ and the decomposition is trivial. □

□

Corollary 21. If q is not prime, P_q has two complete decomposition chains of different length.

As there is a bijection between the subgroups of Γ_0 and the components of $(x^q - x)^{q-1}$ on the right, we will study those subgroups in order to determine whether this polynomial has complete decompositions of different length when q is prime.

Definition 22. $H_0 = \{x + b : b \in \mathbb{F}_q\}$.

Lemma 23. Γ_0 is the semidirect product of H_0 and $\{ax : a \in \mathbb{F}_q^*\}$.

Let G be a subgroup of Γ_0 . As H_0 has prime order, we have two cases:

- $G \cap H_0 = H_0$. Then $H_0 \subseteq G$. If $ax + b \in G$, then for every $b' \in \mathbb{F}_q$ we have $ax + b' \in G$. In particular, $ax \in G$, and $G_0 = \{a \in \mathbb{F}_q^* : ax \in G\} < \mathbb{F}_q^*$. But $\mathbb{F}_q^*$ is cyclic of order $q - 1$, thus G_0 is cyclic of order $m \mid q - 1$. In this case, $G = H_0 \rtimes G_0 \cong C_q \rtimes C_m$.

- $G \cap H_0 = \{x\}$. Then for every $a \in G_0$ there exists exactly one $b \in \mathbb{F}_q$ such that $ax + b \in G$, because $(ax + b) \circ (ax + b')^{-1} = x - b' + b$. As G_0 is cyclic, we have that G is generated by some $a_0x + b_0$ where a_0 generates G_0 and $b_0 \in \mathbb{F}_q$.

This allows to prove the following theorem.

Theorem 24. *If q is prime, then all the maximal chains of subgroups of $\Gamma_0(\mathbb{F}_q)$ have the same length.*

Proof. Let $G_0 = \{x\} < G_1 < \dots < G_n = \Gamma_0(\mathbb{F}_q)$ be a maximal chain. Let $i \in \{1, \dots, n\}$ be such that $G_{i-1} \cap H_0 = \{x\}$ and for all $j \geq i$, $H_0 \subseteq G_j$. For each $j \geq i$ there exists a cyclic group C_i of order m_i with $m_i \mid q - 1$ such that $G_i = H_0 \rtimes C_i$. Thus, the numbers $m_i, m_{i+1}, \dots, m_n$ are a maximal chain of divisors of $q - 1$ greater or equal than m_i .

On the other hand, G_{i-1} must be a cyclic group of order m_i , therefore the orders of $G_1, \dots, G_{i-1}$ are a maximal chain of divisors of m_i .

Therefore, the length of the chain $G_0, \dots, G_n$ is equal to the number of prime factors in a complete factorization of $q - 1$ plus two. $\square \quad \square$

Corollary 25. The polynomial $(x^q - x)^{q-1} \in \mathbb{F}_q[x]$ has maximal decomposition chains of different lengths iff q is not prime.

Remark 26. It is possible to determine all the subgroups of $\Gamma(\mathbb{F}_q)$ by finding all subgroups of $GL(2, q)$. Then all chains of subgroups can be computed, finding out whether the function f has decompositions of different lengths.

4 Conclusions

The results in the last section show some new information about the structure of decompositions of rational functions in the finite case; it is our hope that more can be said about possible versions of Ritt's theorems for finite fields. Also, the algorithms presented here indicate that fast decomposition algorithms in the finite case can be achievable, by using this structure.

References

- [1] C. Alonso, J. Gutierrez, T. Recio, *A rational function decomposition algorithm by near-separated polynomials*. J. Symbolic Comput. 19 (1995), no. 6, 527–544.

- [2] F. Binder, *Polynomial decomposition*. Master's thesis, University of Linz, June 1995.
- [3] C. Corrales-Rodríguez, *A note on Ritt's theorem on decomposition of polynomials*. J. Pure Appl. Algebra 95 (1990), 293–296.
- [4] F. Dorey, G. Whaples, *Prime and composite polynomials*. J. Algebra 28 (1974), 88–101.
- [5] H. T. Engström, *Polynomial substitutions*. Amer. J. Math. 63 (1941), 249–255.
- [6] M. Fried, R. Mac Rae, *On the invariance of chains of fields*. Illinois J. Math. 13 (1969), 165–171.
- [7] S. Lang, *Algebra*. Addison–Wesley, Reading, Mass (1967).
- [8] R. Lidl and H. Niederreiter, *Finite fields*, Cambridge University Press, Cambridge, 1997.
- [9] P. Lüroth, *Beweis eines Satzes über rationale curven*. Math. Ann. 9 (1876), 163–165.
- [10] E. Netto, *Über einen Lüroth-Gordaschen Satz*. Math. Ann. 9 (1895), 310–318.
- [11] J. F. Ritt, *Prime and composite polynomials*. Trans. Amer. Math. Soc. 23 (1922), no. 1, 51–66.
- [12] A. Schinzel, *Selected topics on polynomials*. Ann Arbor, University of Michigan Press, 1982.
- [13] T. W. Sederberg, *Improperly parametrized rational curves*. Computer Aided Geometric Design 3 (1986), 67–75.
- [14] D. Sevilla, *Ritt's Theorems and computation of unirational fields*. Ph. D. Thesis Dissertation. University of Cantabria, 2004.
- [15] E. Steinitz, *Algebraische Theorie der Körper*. Journal für reine und angewante Mathematik, 137 (1910), 167–309.
- [16] B. L. van der Waerden, *Modern Algebra*. Frederick Ungar Publishing Co., New York, 1964.

- [17] R. Zippel, *Rational function decomposition*. Proceedings of ISSAC'91, ACM Press, 1991.

Additional references

- [18] U. Zannier, *Ritt's second theorem in arbitrary characteristic*, Journal reine angew. Math., 445 (1993), 175–203.
- [19] A. Schinzel, *Polynomials with Special Regard to Reducibility*. With an appendix by Umberto Zannier. Encyclopedia of Mathematics and its Applications, 77, Cambridge University Press, Cambridge, 2000.